

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number
WO 01/00665 A2

- (51) International Patent Classification⁷: C07K 14/00
- (21) International Application Number: PCT/US00/17742
- (22) International Filing Date: 27 June 2000 (27.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
- | | | |
|------------|-------------------------------|----|
| 60/141,363 | 28 June 1999 (28.06.1999) | US |
| 60/168,060 | 30 November 1999 (30.11.1999) | US |
| 60/177,836 | 25 January 2000 (25.01.2000) | US |
| 60/178,368 | 27 January 2000 (27.01.2000) | US |
| 60/210,292 | 8 June 2000 (08.06.2000) | US |
- (71) Applicants: OKLAHOMA MEDICAL RESEARCH FOUNDATION [US/US]; 825 N.E. Thirteenth Street, Oklahoma City, OK 73104 (US). THE BOARD OF TRUSTEES OF THE UNIVERSITY OF ILLINOIS [US/US]; 352 Henry Administration Building, 201 S. Wright Street, Urbana, IL 61801 (US).
- (72) Inventors: TANG, Jordan, J., N.; 1204 Leawood Drive, Edmond, OK 73034 (US). HONG, Ling; 7503 Sacramento Drive, Oklahoma City, OK 73139 (US). GHOSH, Arun, K.; 1407 Clinton Place, River Forest, IL 60305 (US).
- (74) Agent: PABST, Patrea, L.; Arnall Golden & Gregory, LLP, 2800 One Atlantic Center, 1201 West Peachtree Street, Atlanta, GA 30309-3450 (US).
- (81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— Without international search report and to be republished upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: INHIBITORS OF MEMAP SIN 2 AND USE THEREOF

WO 01/00665 A2

(57) Abstract: Methods for the production of purified, catalytically active, recombinant memapsin 2 have been developed. The substrate and subsite specificity of the catalytically active enzyme have been determined. The substrate and subsite specificity information was used to design substrate analogs of the natural memapsin 2 substrate that can inhibit the function of memapsin 2. The substrate analogs are based on peptide sequences, shown to be related to the natural peptide substrates for memapsin 2. The substrate analogs contain at least one analog of an amide bond which is not capable of being cleaved by memapsin 2. Processes for the synthesis of two substrate analogues including isosteres at the sites of the critical amino acid residues were developed and the substrate analogues, OMR99-1 and OM99-2, were synthesized. OM99-2 is based on an octapeptide Glu-Val-Asn-Leu-Ala-Ala-Glu-Phe (SEQ ID NO:28) with the Leu-Ala peptide bond substituted by a transition-state isostere hydroxyethylene group (Figure 1). The inhibition constant of OM99-2 is 1.6×10^{-9} M against recombinant pro-memapsin 2. Crystallography of memapsin 2 bound to this inhibitor was used to determine the three dimensional structure of the protein, as well as the importance of the various residues in binding. This information can be used by those skilled in the art to design new inhibitors, using commercially available software programs and techniques familiar to those in organic chemistry and enzymology, to design new inhibitors to memapsin 2, useful in diagnostics and for the treatment and/or prevention of Alzheimer's disease.

INHIBITORS OF MEMAP SIN 2 AND USE THEREOF

Background of the Invention

This invention is in the area of the design and synthesis of specific inhibitors of the aspartic protease Memapsin 2 (beta-secretase) which are useful in the treatment and/or prevention of Alzheimer's Disease.

Alzheimer's disease (AD) is a degenerative disorder of the brain first described by Alois Alzheimer in 1907 after examining one of his patients who suffered drastic reduction in cognitive abilities and had generalized dementia (*The early story of Alzheimer's Disease*, edited by Bick *et al.* (Raven Press, New York 1987)). It is the leading cause of dementia in elderly persons. AD patients have increased problems with memory loss and intellectual functions which progress to the point where they cannot function as normal individuals. With the loss of intellectual skills the patients exhibit personality changes, socially inappropriate actions and schizophrenia (*A Guide to the Understanding of Alzheimer's Disease and Related Disorders*, edited by Jorm (New York University Press, New York 1987). AD is devastating for both victims and their families, for there is no effective palliative or preventive treatment for the inevitable neurodegeneration.

AD is associated with neuritic plaques measuring up to 200 μm in diameter in the cortex, hippocampus, subiculum, hippocampal gyrus, and amygdala. One of the principal constituents of neuritic plaques is amyloid, which is stained by Congo Red (Fisher (1983); Kelly Microbiol. Sci. 1(9):214-219 (1984)). Amyloid plaques stained by Congo Red are extracellular, pink or rust-colored in bright field, and birefringent in polarized light. The plaques are composed of polypeptide fibrils and are often present around blood vessels, reducing blood supply to various neurons in the brain.

Various factors such as genetic predisposition, infectious agents, toxins, metals, and head trauma have all been suggested as possible mechanisms of AD neuropathy. Available evidence strongly indicates that there are distinct types of genetic predispositions for AD. First, molecular analysis has provided evidence for mutations in the amyloid precursor